

# After a video capsule endoscopy with or without a gastroscopy: information for families

---

**This information sheet from Great Ormond Street Hospital (GOSH) describes how to look after your child at home after a video capsule endoscopy with or without a gastroscopy and how to deal with any problems that might occur. If you have any questions, please contact the Gastroenterology Investigation Suite on . Out of hours or in an emergency, please contact your family doctor (GP) or take your child to the nearest Accident and Emergency (A&E) department.**

The test finishes when the video capsule has reached the small intestine. The nurse will remove the recorder and you can take your child home.

## **If the capsule was placed with a gastroscopy under general anaesthetic**

After a general anaesthetic, your child may feel drowsy, light headed or dizzy, and sick or be sick (nausea and vomiting). They may also have itchy skin, a sore throat and bloating.

- You should encourage your child to drink but do not force them to do so. As long as your child is drinking, it does not matter if they do not feel like eating for the first couple of days.
- Your child may be tired and a little clumsy for 24 hours after the anaesthetic, so do not let him or her do anything that might lead to a fall. You should keep your child away from school or nursery for one day until they have recovered from the anaesthetic.

## **Passing the capsule**

Your child should pass the video capsule with 24 to 48 hours of the test stopping. You may be able to spot it in your child's faeces (poo) – it is disposable so can be flushed away in the toilet. The capsule may be inside a lump of faeces so you may not be able to see it – please do not be concerned.

However, please seek medical advice if your child:

- Has tummy pain or bloating that is unusual for them or worse than usual
- They vomit large amounts of bright red blood
- They become generally unwell, refusing to eat or drink or very sleepy

## **Getting back to normal**

Unless you have been advised otherwise

- Your child can start eating and drinking their usual diet
- They can re-start any medications that were stopped for the test

### **Getting the test results**

The results of the test will usually be reviewed and available with 10 working days. If you do not have a clinic appointment arranged with the consultant who referred your child for the test, please arrange one. Their number to call will be on the bottom of your last appointment letter.

Ref: 2017F2091 March 2018

Compiled by the Gastroenterology Investigation Suite in collaboration with the Child and Family Information Group