


# Tetralogy of Fallot (TOF)


## With Hypoplastic Pulmonary Annulus and Pulmonary Artery



To correct the problem, the hole between the bottom chambers is closed. This is done using a patch of a material called Dacron®. Also, because the hole is closed, the narrowing under the pulmonary valve is relieved. This allows blood from the bottom right heart chamber to flow into the lungs as it should.

When the pulmonary annulus and pulmonary artery are underdeveloped, the correction involves relieving the narrowings. This is done with an incision across the pulmonary annulus and artery and enlarging it with a wide rectangular-shaped patch. The patch is a piece of the sac that the heart sits in. This sac is called pericardium.

The surgery is done through a median sternotomy (chest) incision. If a shunt was placed earlier, it is also removed.




There are 4 heart problems:

1. a hole between the bottom 2 chambers of the heart (ventricular septal defect)
2. a narrowing under or at the pulmonary valve (pulmonary stenosis)

There is also underdeveloped (hypoplastic) pulmonary annulus (ring-shaped tissue of the pulmonary artery). The pulmonary artery is also narrowed.

3. enlargement of the bottom right side of the heart
4. aorta is lined up just over the hole between the bottom 2 chambers

Children with TOF have mixing of the red and blue blood through the hole between the bottom two chambers of the heart. Blood going to the lungs is restricted by the narrowing under the pulmonary valve. This protects the lungs from getting too much blood flow. The bottom right side of the heart becomes enlarged. This happens because it must hold too much blood and try to squeeze it through the narrowing to get to the lungs for oxygen.



Sometimes an infant is too sick or the pulmonary arteries are too small for the corrective surgery. In that case, a small tube (either from a blood vessel of the infant or artificial material) is placed from the subclavian artery into the pulmonary artery (Blalock Taussig Shunt). The tube (shunt) allows blood to enter the lungs at all times. The shunt is placed through a thoracotomy (side) incision. The shunt is removed at the time of the correction.